

On the dimension of splines spaces over T-meshes with smoothing cofactor-conformality method

X. Li

Department of Mathematics, USTC,
Hefei, Anhui Province 230026, P. R. China.
lixustc@ustc.edu.cn

October 22, 2012

Abstract

The present paper provides a general formula for the dimension of spline space over T-meshes using smoothing cofactor-conformality method. And we introduce a new notion, **Diagonalizable T-mesh**, over which the dimension formula is only associated with the topological information of the T-mesh. A necessary and sufficient condition for characterization a diagonalizable T-mesh is also provided. Using this technique, we find that the dimension is possible instable under the condition of [1] and we also provide a new correction theorem.

Keywords: Dimension, T-splines, Spline space, T-mesh, smoothing cofactor-conformality method

1 Introduction

NURBS (Non-Uniform Rational B-Spline) is the standard for generating and representing free-form curves and surfaces, which is a basic tool for using in CAD [2] and is also a desirable tool for iso-geometric analysis [3]. A well-known and significant disadvantage of NURBS is that it is based on a tensor product structure with a global knot insertion operation. It is desirable to generalize NURBS to spline space which can handle hanging nodes.

Spline spaces over T-meshes $\mathcal{S}(d_1, d_2, \alpha, \beta, \mathcal{T})$ were first introduced in [4], which is a bi-degree (d_1, d_2) piecewise polynomial spline space over T-mesh $\mathcal{T}$ with smoothness order α and β in two directions. Spline space over T-meshes have been applied in fitting [5], stitching [6], simplification [7], isogeometric analysis [8], [9], solving elliptic equations [10] and spline space over triangulations with hanging nodes [11]. Spline over T-mesh has several advantages. For example, it has a simple local refinement which will never introduce additional refinement according to the definition. And it is a polynomial in each face which has simple and efficient integration for numerical analysis.

NURBS with local refinement is discovered before spline over T-mesh, called "T-spline", which are defined on a T-mesh by certain collections of B-splines functions defined on the mesh [12], [13]. T-splines have been proved to be a powerful free-form geometric shape technology that solves most of the limitations inherent in NURBS. T-splines have several advantages over NURBS such as local refinement [13] and watertightness [14], and they are forward and backward compatible with NURBS. These capabilities make T-splines attractive both for CAD and also desirable for iso-geometric analysis [15]. However, it is very difficult to unravel the mystery of T-spline spaces which have important implications in establishing approximation, stability, and error estimates [16]. And till now, only a sub-class of T-spline space, analysis-suitable T-splines space [17], [18], [19] has been discovered using the technology from spline space over T-meshes.

Thus, it is very important to understand the spline space of piecewise polynomials of a given smoothness on a T-mesh, which foundation but non-trivial step is to calculate the dimension of the space. Till now, many different methods have been applied to tackle these issues, such as B-net [4], minimal determining set [20], smoothing cofactor-conformality method [1] and homological technique [21]. B-net and minimal determining set methods are suitable for spline space with reduced regularity, i.e., the degrees for the polynomial is larger enough than the smoothness order. Smooth cofactor-conformality method [22], [23] is a power tool for calculating the dimension of spline space in multi-variate splines theory. It transfers the smoothness conditions into algebraic forms and calculate the dimension using linear algebraic tools. Homological technique is another power tool for calculating the dimension using the similar idea except regarding the smoothness conditions as the kernel of a certain linear maps. The present paper is focusing on smoothing cofactor-conformality method.

All the existing dimension results are forcing the T-mesh to be special nesting structure, such as hierarchical, regular T-subdivision and no cycles [20]. With such structure, one can calculate the dimension level by level. Our approach is based on a different point of view. As smoothing cofactor-conformality method can convert the smoothness conditions into algebraic forms, so we directly focus on the algebraic forms and study the condition, under which the dimension of the linear system doesn't associated with knot values. And then we use the condition to find a new notion, diagonalizable T-meshes, is the corresponding T-meshes. We believe that this new notion is the key condition to compute the dimension using smoothing cofactor-conformality method. The main contribution of the present paper includes,

- We provide a general formula for the dimension of spline space over any regular T-meshes without holes using smoothing cofactor-conformality method.
- We provide a new notion, diagonalizable T-mesh, over which the dimension formula is only associated with the topological information of the T-mesh. We also provide a necessary and sufficient condition for characterization diagonalizable T-meshes;

- We discover that the dimension is possible instable under the condition of [1] and we also provide a new correction theorem.

The remaining paper is structured as follows. Pertinent background on spline space over T-mesh is reviewed in Section 2. In Section 3, we review how to use smoothing cofactor-conformality method to analysis the dimension of spline space over T-meshes. In section 4, we provide a condition under which we can calculate the dimension of spline space over T-mesh without considering the knot values. In section 5, we first show that the dimension result in [1] is not correct and we provide a new correction based on this new technology. The last section is conclusion and future work.

2 T-mesh and spline over T-mesh

In this section, we briefly review the notion of spline spaces over T-meshes and the related dimension results of the corresponding spline space.

2.1 T-mesh

A T-mesh $\mathcal{T}$ is a collection of axis-aligned rectangles F_i such that the interior of the domain Ω is $\cup F_i$, and the distinct rectangles F_i and F_j can only intersect at points on their edges. The rectangles F_i are also called the *face* or *cell* of the T-mesh. The vertices of the rectangles are called the *nodes* or *vertices* for a T-mesh. The line segment connecting two adjacent vertices on a grid line is called an *edge* of the T-mesh. T-meshes include tensor-product meshes as a special case. However, in contrast to tensor-product meshes, T-meshes are allowed to have *T-junctions*, or *T-nodes*, which are vertices of one rectangle that lies in the interior of an edge of another rectangle. The domain Ω need not be rectangular, which may have holes, concave corners. For example the T-mesh in Figure 1, the grey region is a hole and vertices V_{38} and V_{88} are both concave corners. In the present paper, we require the T-meshes to be **regular** and **without holes**. Here regular means that the set of all rectangles for a T-mesh containing a vertex has a connected interior [24].

The vertices, edges can be divided into two parts. If a vertex is on the boundary grid line of the T-mesh, then is called a *boundary vertex*. Otherwise, it is called an *interior vertex*. If both vertices of an edge are boundary vertices, then it is called a *boundary edge*; otherwise it is called an *interior edge*. An *l-edge* is a line segment which consists of several interior edges. It is the longest possible line segment, which interior edges are connected and two end points being T-junctions or boundary vertices. l-edges have three different classes. If the two end vertices of a l-edge are interior vertices, then the l-edge is called *interior l-edge*. If two end vertices of a l-edge are both boundary vertices, then the l-edge is called a *cross-cut*. Otherwise, if one end vertex is boundary vertex and the other is interior vertex, then the l-edge is called a *ray*. A *mono-vertex* is the intersection vertex of an interior l-edge and a cross-cut or a ray and a *free-vertex* is the intersection between cross-cuts and rays. For example, in Figure 1,




Figure 1: A T-mesh.

vertices V_{53} , V_{44} and V_{46} are interior vertices, and V_{15} , V_{16} and V_{95} are boundary vertices. The l-edge $V_{15}V_{95}$ is a cross-cut, while $V_{16}V_{46}$ is a ray, and $V_{44}V_{84}$ and $V_{57}V_{58}$ are interior l-edges.

For later use we introduce some notations for a T-mesh as shown in Table 1

2.2 Spline space over T-mesh

Given a T-mesh $\mathcal{T} \in \mathbb{R}^2$, let $\mathcal{F}$ denote all the cells in $\mathcal{T}$ and Ω the region occupied by all the cells in $\mathcal{T}$. The bi-degree (d_1, d_2) polynomial spline space over T-mesh $\mathcal{T}$ with smoothness order α and β is defined as

$$\mathcal{S}(d_1, d_2, \alpha, \beta, \mathcal{T}) := \left\{ f(x, y) \in C^{\alpha, \beta}(\Omega) \mid f|_{\phi} \in P_{d_1 d_2}, \forall \phi \in \mathcal{F} \right\},$$

where $P_{d_1 d_2}$ is the space of all the polynomials with bi-degree (d_1, d_2) , and $C^{\alpha, \beta}(\Omega)$ is the space consisting of all the bivariate functions which are continuous in Ω with order α along x direction and with order β along y direction. It is obvious that $\mathcal{S}(d_1, d_2, \alpha, \beta, \mathcal{T})$ is a linear space, which is called the spline space over the given T-mesh $\mathcal{T}$.

Until now, several articles have been studied to analysis the dimension of the spline space over some special families of T-meshes.

- **Reduced regularity:**

In 2006, [4] studied the dimension of the spline space under the constrains that the order of the smoothness is less than half of the degree of the spline functions. According to Theorem 4.2 in [4], it follows that if $d_1 \geq 2\alpha + 1$ and $d_2 \geq 2\beta + 1$,

$$\dim \mathcal{S}(d_1, d_2, \alpha, \beta, \mathcal{T}) = F(d_1 + 1)(d_2 + 1) - E_h(d_1 + 1)(\beta + 1) - E_v(d_2 + 1)(\alpha + 1) + V(\alpha + 1)(\beta + 1),$$

Table 1: Notations for a T-mesh	
F	number of faces in $\mathcal{T}$
E^h	number of horizontal interior edges in $\mathcal{T}$
E^v	number of vertical interior edges in $\mathcal{T}$
V	number of interior vertices in $\mathcal{T}$
C^h	number of horizontal cross-cuts in $\mathcal{T}$
C^v	number of vertical cross-cuts in $\mathcal{T}$
T^h	number of horizontal interior l-edges in $\mathcal{T}$
T^v	number of vertical interior l-edges in $\mathcal{T}$
n_e	number of interior l-edges in $\mathcal{T}$ ($T^h + T^v$)
V^+	number of free-vertices in $\mathcal{T}$
N^h	minimal integer larger or equal to $\frac{d_1+1}{d_1-\alpha}$
N^v	minimal integer larger or equal to $\frac{d_2+1}{d_2-\beta}$
n_c	$(d_1 - \alpha)(d_2 - \beta)(V - V^+)$
n_r	$(d_1 + 1)(d_2 - \beta)T^h + (d_2 + 1)(d_1 - \alpha)T^v$

where F , E_h , E_v , and V are defined in Table 1. [24] also proved this result using minimal determining set method. And later, [25] analysis a special T-spline with reducing regularity using the dimension in [4].

- **Enough mono-vertices**

In 2006, [1] calculated the dimension of spline space over a T-mesh if each interior l-edges have enough mono-vertices. In the T-mesh, if the interior of each horizontal interior l-edge has at least $N^h - 2$ mono-vertices and the interior of each vertical interior edge segment has at least $N^v - 2$ mono-vertices, then the dimension of spline space over the T-mesh is,

$$\dim \mathcal{S}(d_1, d_2, \alpha, \beta, \mathcal{T}) = (d_1 + 1)(d_2 + 1) + (C_h - T_h)(d_1 + 1)(d_2 - \beta) + (C_v - T_v)(d_2 + 1)(d_1 - \alpha) + V(d_1 - \alpha)(d_2 - \beta),$$

here C_h , C_v , T_h , T_v and V are defined in Table 1. However, we will show that the condition in this paper is not right.

- **Instability:**

In 2011, [26, 27] discovered that the dimension of the associated spline space is instability over some particular T-meshes, i.e, the dimension is not only associated with the topological information of the T-mesh but also associated with the geometric information of the T-mesh.

- **Analysis-suitable T-splines:**

In 2011, [17, 18] provided a mildly restricted subset of T-splines, which optimized to meet the needs of both design and analysis. And [19] compute the dimension of the spline space $\mathcal{S}(d, d, d - 1, d - 1, \mathcal{T})$ if the T-mesh $\mathcal{T}$ is an extended T-mesh of an analysis-suitable T-mesh.

- **Regular T-subdivision:**

[21] studied the dimension for spline space $\mathcal{S}(d_1, d_2, \alpha, \beta, \mathcal{T})$ when the

T-mesh is a regular T-subdivision by exploiting homological techniques, which is a special case of the present paper.

- **Special hierarchical T-mesh:**

[28] provided the dimension for spline space $\mathcal{S}(d, d, d-1, d-1, \mathcal{T})$ over a special hierarchical T-mesh using homological algebra technique.

3 Smoothing cofactor-conformality method

In this section, we will review one of the main methods, smooth cofactor-conformality method introduced in [22] and [23], for computing the dimension of spline space over T-meshes.

In the theory of multi-variate splines, in order to calculate the dimension of a spline space, one first needs to transfer the smoothness conditions into algebraic forms.

Referring to Figure 2, for any interior vertex $v_{i,j} = (x_i, y_j)$, suppose the four surrounding bi-degree (d_1, d_2) polynomial patches are $p_{i,j}^k(x, y), k = 0, 1, \dots, 3$ respectively (if the vertex $v_{i,j}$ is a T-junction, then some of the polynomial patches are identical). For example for the left T-junction in Figure 2 b, patches $p_{i,j}^1(x, y), p_{i,j}^2(x, y)$ are identical.




Figure 2: smoothing cofactors around a vertex.

As $p_{i,j}^0(x, y)$ and $p_{i,j}^1(x, y)$ are C^α continuity, so there exists a bi-degree $(d_1 - \alpha - 1, d_2)$ polynomial $\lambda_{i,j}^2(y)$ such that

$$p_{i,j}^1(x, y) - p_{i,j}^0(x, y) = \lambda_{i,j}^2(x, y)(x - x_i)^{\alpha+1}, \quad (1)$$

Here $\lambda_{i,j}^2(x, y)$ is called **edge cofactor** for the common edge of patches $p_{i,j}^0(x, y)$ and $p_{i,j}^1(x, y)$. If two patches are identical, then the edge cofactor is zero.

Similarly, there also exist bi-degree $(d_1 - \alpha - 1, d_2)$ polynomial $\lambda_{i,j}^1(x, y)$,

bi-degree $(d_1, d_2 - \beta - 1)$ polynomials $\mu_{i,j}^1(x, y)$ and $\mu_{i,j}^2(x, y)$, such that

$$p_{i,j}^2(x, y) - p_{i,j}^1(x, y) = \mu_{i,j}^1(x, y)(y - y_j)^{\beta+1}, \quad (2)$$

$$p_{i,j}^3(x, y) - p_{i,j}^2(x, y) = -\lambda_{i,j}^1(x, y)(x - x_i)^{\alpha+1}, \quad (3)$$

$$p_{i,j}^0(x, y) - p_{i,j}^3(x, y) = -\mu_{i,j}^2(x, y)(y - y_j)^{\beta+1}. \quad (4)$$

Sum with all these equations, we have

$$(\lambda_{i,j}^1(x, y) - \lambda_{i,j}^2(x, y))(x - x_i)^{\alpha+1} = (\mu_{i,j}^1(x, y) - \mu_{i,j}^2(x, y))(y - y_j)^{\beta+1}. \quad (5)$$

Since $(x - x_i)^{\alpha+1}$ and $(y - y_j)^{\beta+1}$ are prime to each other, so there exist bi-degree $(d_1 - \alpha - 1, d_2 - \beta - 1)$ polynomial $d_{i,j}(x, y)$, such that,

$$\lambda_{i,j}^1(x, y) - \lambda_{i,j}^2(x, y) = d_{i,j}(x, y)(y - y_j)^{\beta+1}, \quad \mu_{i,j}^1(x, y) - \mu_{i,j}^2(x, y) = d_{i,j}(x, y)(x - x_i)^{\alpha+1}.$$

Let

$$d_{i,j}(x, y) = \sum_{p=0}^{d_1-\alpha-1} \sum_{q=0}^{d_2-\beta-1} d_{i,j}^{p,q} (x - x_i)^p (y - y_j)^q,$$

We call these $d_{i,j}^{p,q}$ **vertex cofactor**. Denote $\hat{\mathbf{d}}_{i,j}$ to be a vector contains all coefficients $d_{i,j}^{p,q}$, i.e.,

$$\hat{\mathbf{d}}_{i,j} = (d_{i,j}^{0,0}, d_{i,j}^{0,1}, \dots, d_{i,j}^{d_1-\alpha-1, d_2-\beta-2}, d_{i,j}^{d_1-\alpha-1, d_2-\beta-1}).$$

For boundary vertices, there is a little different. Since in this case, we only have parts of the four equations such as (1),(2),(3),(4). So we don't need to assign the bi-degree $(d_1 - \alpha - 1, d_2 - \beta - 1)$ polynomial $d_{i,j}(x, y)$ for the boundary vertex. Instead, we will assign the edge cofactor for the corresponding edge. For example, in Figure 3a, we need two edge cofactors $\lambda_{i,j}^1$ and $\mu_{i,j}^1$ and for the figure b, we need only one edge cofactors $\mu_{i,j}^1$.





Figure 3: smoothing cofactors around a boundary vertex. The grey regions are outside of the T-mesh.



(a) a.

Figure 4: smoothing cofactors along a horizontal edge segment.

The vertex cofactors for the interior vertices are not totally free, there are other constrains for the continuity condition along each l-edge in the T-mesh.

We first consider a horizontal *interior l-edge* referring to Figure 4 with $k+1$ vertices $v_{i_t,j}$, $t = 0, 1, \dots, k$. According to equation (6), we have

$$\begin{aligned}
 \mu_{i_0,j}^1(x, y) - 0 &= d_{i_0,j}(x, y)(x - x_{i_0})^{\alpha+1}, \\
 &\dots \\
 \mu_{i_t,j}^1(x, y) - \mu_{i_t,j}^2(x, y) &= d_{i_t,j}(x, y)(x - x_{i_t})^{\alpha+1}, \\
 &\dots \\
 0 - \mu_{i_k,j}^2(x, y) &= d_{i_k,j}(x, y)(x - x_{i_k})^{\alpha+1}.
 \end{aligned} \tag{6}$$

and

$$\mu_{i_{t-1},j}^2(x, y) = \mu_{i_t,j}^1(x, y).$$

Sum all these equations, we have the following equation,

$$\sum_{t=0}^k d_{i_t,j}(x, y)(x - x_{i_t})^{\alpha+1} = 0. \tag{7}$$

Similarly, the constrains for a vertical interior l-edge is

$$\sum_{t=0}^l d_{i,j_t}(x, y)(y - y_{j_t})^{\beta+1} = 0. \tag{8}$$

The above equations are called **edge conformality conditions**.

Lemma 3.1. Assume each x_{i_t} and y_{j_t} are distinct, then the dimensions of solution space of equation (7) and (8) are $(d_2 - \beta)(l(d_1 - \alpha) - d_1 - 1)_+$ and $(d_1 - \alpha)(k(d_2 - \beta) - d_2 - 1)_+$ respectively.

Proof. See [1] for more details. $\square$

Remark 3.2. If the number of the vertices in a horizontal l-edge is less than N^h , or the number of the vertices in a vertical l-edge is less than N^v , then the l-edge will not contribute the dimension of the spline space, i.e., we can delete the l-edge without altering the spline space. Thus, we called such l-edge **vanished l-edge**. In the following, we assume that all l-edges are not vanished l-edges.

Now we consider a horizontal ray with $r + 1$ vertices $\nu_{i_t,j}, t = 0, 1, \dots, k$. Without loss generalization, we assume $\nu_{i_0,j}$ is a boundary vertex. According to equation (6), we have

$$\begin{aligned} \mu_{i_1,j}^1(x, y) - \mu_{i_1,j}^2(x, y) &= d_{i_1,j}(x, y)(x - x_{i_1})^{\alpha+1}, \\ &\dots \\ \mu_{i_t,j}^1(x, y) - \mu_{i_t,j}^2(x, y) &= d_{i_t,j}(x, y)(x - x_{i_t})^{\alpha+1}, \\ &\dots \\ 0 - \mu_{i_k,j}^2(x, y) &= d_{i_k,j}(x, y)(x - x_{i_k})^{\alpha+1}. \end{aligned} \quad (9)$$

and

$$\mu_{i_{t-1},j}^2(x, y) = \mu_{i_t,j}^1(x, y).$$

Sum all these equations, we have the following equation,

$$\sum_{t=1}^k d_{i_t,j}(x, y)(x - x_{i_t})^{\alpha+1} = \mu_{i_1,j}^2(x, y). \quad (10)$$

The main different of the constraint from (7) and (8) is that these is one edge cofactor $\mu_{i_1,j}^2(x, y)$ in the constraints. For any $d_{i_t,j}(x, y)$ assigned for the interior vertices, we can assign $\mu_{i_1,j}^2(x, y)$ as $\sum_{t=1}^k d_{i_t,j}(x, y)(x - x_{i_t})^{\alpha+1}$ to satisfy the constraint.

For a horizontal cross-cut in the T-mesh, since it has two boundary vertices, we can conclude that it has $(d_1 + 1)(d_2 - \beta)$ degrees of freedoms. Similarly, a vertical cross-cut has $(d_2 + 1)(d_1 - \alpha)$ degrees of freedoms.

The linear systems as (7) or (8), associated with each interior l-edges can be assembled into a global system as $\mathcal{M}\mathbf{x} = 0$. Here $\mathcal{M}$ is a $n_r \times n_c$ matrix, which is called **conformality conditions matrix**. And $\mathbf{x}$ is a column vector whose elements are all the vertex cofactors for the interior vertices in the T-mesh.

And according to the above anlysis, we can get the dimension for spline space over any general T-mesh without holes according to smoothing cofactor-conformality method which is stated in the following theorem.

Theorem 3.3. *Given a T-mesh $\mathcal{T}$ which has no vanished l-edges and holes, let matrix $\mathcal{M}$ be the conformality conditions matrix, then the dimension of spline space over the T-mesh is,*

$$\begin{aligned} \dim \mathcal{S}(d_1, d_2, \alpha, \beta, \mathcal{T}) &= (d_1 + 1)(d_2 + 1) + C^h(d_1 + 1)(d_2 - \beta) + \\ &\quad C^v(d_2 + 1)(d_1 - \alpha) + V^+(d_1 - \alpha)(d_2 - \beta) + \dim \mathcal{M}, \end{aligned}$$

4 Diagonalizable

Theorem 3.3 indicates that the main difficult to compute the dimension of spline space over T-mesh is to calculate the rank of conformality condition matrix $\mathcal{M}$. It is obvious that the structure of the matrix is associated with the order of

edge conformality conditions and it is also associated with the order of vertices cofactors. Most of existing method study the dimension of the matrix by forcing the T-mesh to be nest structure, such as T-subdivision, hierarchical. However, nesting structure should be fine for application, but it is hiding some essential properties for T-mesh. What we wish to answer is that under what condition can we compute the dimension regardless the knot intervals for a given T-mesh and why such condition is important.

Definition 4.1. *Given a T-mesh $\mathcal{T}$, suppose we order the all the interior l-edges as $e_{i_j}, j = 1, 2, \dots, n_e$, then we can compute the **new-vertex-vector** ν^i . Here ν_1^i is the number of vertices on l-edge e_{i_1} and ν_j^i is the number of vertices on l-edge e_{i_j} but not on l-edges $e_{i_k}, k < j$.*

Definition 4.2. *A T-mesh is called **Diagonalizable** if there exists an order of l-edges $e_{i_j}, j = 1, 2, \dots, n_e$ such that the new-vertex-vector ν^i satisfies that $\nu_j^i \geq N^h$ if e_{i_j} is horizontal and $\nu_j^i \geq N^v$ if e_{i_j} is vertical.*

Lemma 4.3. *If a T-mesh is diagonalizable, then the matrix $\mathcal{M}$ has full column rank regardless the knot intervals.*

Proof. Since we assume that the T-mesh has no vanished l-edges, the matrix $\mathcal{M}$ has more columns than rows, i.e., $n_c \leq n_r$. After arranging the order of edge conformality conditions and the order of vertex cofactors, an appropriate partition of the linear system of constraints is

$$\left[\begin{array}{c|c} \mathcal{M}_1 & \mathcal{M}_2 \end{array} \right] \begin{bmatrix} \mathbf{x}_1 \\ \mathbf{x}_2 \end{bmatrix} = \mathbf{0} \quad (11)$$

where $\mathcal{M}_1$ is a $n_r \times n_r$ matrix and $\mathcal{M}_2$ is a $n_r \times (n_c - n_r)$ matrix, $\mathbf{x}_1^T$ is a vector of the first n_r vertex cofactors, and $\mathbf{x}_2$ is a vector of the remaining vertex cofactors.

Since the T-mesh is diagonalizable, so there exist the order for the interior l-edges satisfy the condition stated in definition 4.2. Without loss of generalization, we assume the order for the l-edges are $e_i, i = 1, 2, \dots, n_e$. Then we arrange the order of the edge conformality conditions corresponding to l-edge from e_{n_e} to e_1 . The order of the vertex cofactors can be placed in the following fashion. For l-edge e_{n_e} , if it is a horizontal l-edge, then there exist N^h vertices which are not appeared in the other l-edges. Each vertex corresponds $(d_1 - \alpha) \times (d_2 - \beta)$ cofactors, so these N^h vertices correspond $(d_1 - \alpha) \times (d_2 - \beta) N^h \geq (d_1 + 1)(d_2 - \beta)$ cofactors. Select any $(d_1 + 1)(d_2 - \beta)$ cofactors and put in the beginning of $\mathbf{x}_1^T$. If the l-edge is a vertical l-edge, we will select $(d_2 + 1)(d_1 - \alpha)$ cofactors and put them in the beginning of $\mathbf{x}_1^T$. These process can be applied for the remaining l-edges. And then the matrix $\mathcal{M}_1$ is in upper block triangular form and according to Lemma 3.1 each diagonal block $(d_2 - \beta)(d_1 + 1) \times (d_2 - \beta)(d_1 + 1)$ or $(d_1 - \alpha)(d_2 + 1) \times (d_1 - \alpha)(d_2 + 1)$ matrix is full rank, thus matrix $\mathcal{M}_1$ is obviously of full rank. $\square$

Theorem 4.4. Suppose T -mesh $\mathcal{T}$ is diagonalizable and has no vanished l -edges and holes, then the dimension of spline space over the T -mesh is,

$$\dim \mathcal{S}(d_1, d_2, \alpha, \beta, \mathcal{T}) = (d_1 + 1)(d_2 + 1) + (C^h - T^h)(d_1 + 1)(d_2 - \beta) + (C^v - T^v)(d_2 + 1)(d_1 - \alpha) + V(d_1 - \alpha)(d_2 - \beta).$$

Proof. As T -mesh $\mathcal{T}$ is diagonalizable, so

$$\dim \mathcal{M} = n_c - n_r = (d_1 - \alpha)(d_2 - \beta)(V - V^+) - (d_1 + 1)(d_2 - \beta)E^h - (d_2 + 1)(d_1 - \alpha)E^v.$$

According to theorem 3.3, we have the dimension $\dim \mathcal{S}(d_1, d_2, \alpha, \beta, \mathcal{T})$ is

$$\begin{aligned} \dim &= (d_1 + 1)(d_2 + 1) + C^h(d_1 + 1)(d_2 - \beta) + C^v(d_2 + 1)(d_1 - \alpha) + V^+(d_1 - \alpha)(d_2 - \beta) + \\ &\quad (d_1 - \alpha)(d_2 - \beta)(V - V^+) - (d_1 + 1)(d_2 - \beta)E^h - (d_2 + 1)(d_1 - \alpha)E^v \\ &= (d_1 + 1)(d_2 + 1) + (C^h - T^h)(d_1 + 1)(d_2 - \beta) + (C^v - T^v)(d_2 + 1)(d_1 - \alpha) + \\ &\quad V(d_1 - \alpha)(d_2 - \beta). \end{aligned}$$

□

Now we provide several examples for dimension of spline space over the following T -meshes.




Figure 5: Two example T -meshes for spline space $\mathcal{S}(3, 3, 2, 2, \mathcal{T})$.

Example 4.1. The first examples are spline space $\mathcal{S}(3, 3, 2, 2, \mathcal{T})$ over the two T -meshes in Figure 5.

The first T -mesh has some concave corners and one interior l -edge which is a vanished l -edge since it has only two vertices. And the T -mesh has two cross-cut. So according to theorem 3.3, the dimension of the spline space $\mathcal{S}(3, 3, 2, 2, \mathcal{T})$ over the first T -mesh is $16 + 2 \times 4 = 24$.

The second T -mesh is much more complex. It has four interior l -edges $e_i, i = 1, \dots, 4$, four cross-cuts and five rays. If we arrange l -edges as e_1, e_2, e_3, e_4 , then the new-vertex-vector corresponds this order is $(5, 5, 4, 3)$. But we

can arrange the l -edges as order e_1, e_4, e_3, e_2 , then the new-vertex-vector corresponds this order is $(5, 4, 4, 4)$, which means the T -mesh is diagonalizable. In the T -mesh, we have 4 cross-cuts, 2 horizontal and 2 vertical interior l -edges and 31 interior vertices, so according to theorem 4.4, the dimension of the spline space $\mathcal{S}(3, 3, 2, 2, \mathcal{T})$ over the first T -mesh is $16 + 4(4 - 4) + 31 = 47$. We should mention that non of existing method can calculate the dimension for this example since the T -mesh is not T -subdivision or hierarchical.




Figure 6: Non-diagonalizable vs. diagonalizable for spline space $\mathcal{S}(3, 3, 1, 1, \mathcal{T})$.

Example 4.2. The second example is associated with spline space $\mathcal{S}(3, 3, 1, 1, \mathcal{T})$ over the two T -meshes in Figure 6.

The first T -mesh has four interior l -edges. And we can see that it is not diagonalizable since the new-vertex-vector could be $(3, 2, 2, 1)$ or $(3, 3, 1, 1)$ for different orders. Thus, we cannot complete the dimension using the current method. Actually, according to our knowledge, no existing method can compute the dimension of such spline space.

The second T -mesh also has four interior l -edges, but according to our knowledge, no existing method can compute the dimension of such spline space. But if we arrange the order of the l -edges to be e_1, e_2, e_3 and e_4 , we can see that the new-vertex-vector is $(3, 2, 2, 2)$, i.e., the T -mesh is diagonalizable. Since the T -mesh has 8 cross-cut, 4 interior l -edges, and 27 interior vertices. So the dimension of the spline space $\mathcal{S}(3, 3, 1, 1, \mathcal{T})$ over the second T -mesh is $16 + 4 \times 2 \times (8 - 4) + 27 \times 4 = 156$.

Example 4.3. The third example is associated with spline space $\mathcal{S}(3, 3, 2, 2, \mathcal{T})$ over the two T -meshes in Figure 7. Both T -meshes have four interior l -edges and we can check that the T -meshes are not diagonalizable. For the first T -mesh, the dimension is 65 and the dimension for the second T -mesh is instable, i.e., the dimension is associated with the value of the knots.

We can see that the second T -mesh can be constructed by moving two red l -edges of the first T -mesh. Although the two T -meshes have different structure, but if we look at the structure of the conformality conditions matrix, we can see that the structure of two matrixes are identical, except the two red l -edges using




Figure 7: The dimension of bi-cubic spline space over the T-mesh is instable, which is generated from the left T-mesh by moving two red l-edges.

the knots of s_2 to s_6 instead of s_1 to s_5 . So diagonalizable is a concept for the structure of the conformality conditions matrix not T-mesh itself.

This example also tells us that if a T-mesh is not diagonalizable, then we have to consider the knot values in order to analysis the dimension of the spline space using smoothing cofactor-conformality method.

Remark 4.5. A similar result has been obtained in [21], which provides the dimension for a special T-mesh, called regular T-subdivision. The main difference between these two method is that the diagonalizable T-meshes don't need to be nested structure. Regular T-subdivision is a special case of diagonalizable T-mesh.

4.1 Characterization

In this section, we will provide a necessary and sufficient condition for characterization a diagonalizable T-mesh.

Lemma 4.6. A necessary and sufficient condition for a T-mesh to be diagonalizable is for any interior l-edges set $\mathbb{S}$, there at least exists one horizontal l-edge such that the number of vertices on this l-edge but not on the other l-edge in $\mathbb{S}$ is at least N^h , or there at least exists one vertical l-edge such that the number of vertices on this l-edge but not on the other l-edge in $\mathbb{S}$ is at least N^v .

Proof. First, we prove the condition is necessary using reduction to absurdity. IF the T-mesh is diagonalizable, but there exists a set of l-edges $\{e_{i_1}, e_{i_2}, \dots, e_{i_s}\}$ such that any horizontal l-edges in the set at most have N^h vertices which are not on the other l-edges in the set and any vertical l-edges at most have N^v vertices which are not on the other l-edges in the set. And since the T-mesh is diagonalizable, so without loss of generalization, we assume when the order

of the interior l-edges is $e_1, e_2, \dots, e_{n_e}$, in which order the l-edges satisfy the condition of diagonalizable. Let k to the maximal index for all $i_j, j = 1, \dots, s$. Now, we consider l-edge e_k , since it has at most $N^h - 1$ or $N^v - 1$ vertices which are not on the other l-edges in the set, so it also has at most $N^h - 1$ or $N^v - 1$ vertices are not on the l-edges for $e_1, \dots, e_{k-1}$ since $\mathbb{S} \subseteq \{e_1, \dots, e_{k-1}, e_k\}$, which violates the assumption of diagonalizable. Thus, the condition is necessary.

Now we prove the condition is sufficient. For the set of l-edges $e_i, i = 1, \dots, n_e$, according to the assumption, there exist one l-edge which has enough vertices on the l-edge but not on the others. Without loss of generalization, we assume it is e_1 . Suppose we have ordered the l-edges as $e_1, e_2, \dots, e_j$ satisfy the diagonalizable condition, then for set $\{e_{j+1}, \dots, e_{n_e}\}$, according to the assumption, there exist one l-edge which has enough vertices on the l-edge but not on the others. Without loss of generalization, we assume it is e_{j+1} . With this process, we can order the l-edges such that it satisfy the diagonalizable condition, which completes the proof. $\square$

5 Correction for [1]

In this section, we will show that the dimension result in [1] is not right. Precisely, the dimension under the condition of [1] is possible instable. And we also provide a new modified theorem using the technology builded in the last section.

5.1 A instable example under the condition of [1]




Figure 8: Counterexample for paper [1].

Consider the spline space $\mathcal{S}(3, 3, 2, 2, \mathcal{T})$ over the T-mesh illustrated in Figure 8. There are four interior l-edges in the T-mesh and each one have two macro-vertices in the interior, which satisfy the condition in [1]. However, we

will analysis the dimension in the following and show that the dimension of the spline space over the T-mesh is instable.

Actually, we arrange the interior l-edges as the order of e_1, e_2, e_3 and e_4 . And we arrange the order of the vertices as $v_{2,2}, v_{3,2}, v_{4,2}, v_{8,2}, v_{7,2}, v_{7,3}, v_{7,4}, v_{7,8}, v_{7,7}, v_{6,7}, v_{5,7}, v_{1,7}, v_{2,7}, v_{2,6}, v_{2,5}, v_{2,1}$ and $v_{2,2}$, then we can get the sparse matrix $\mathcal{M}$ which has the following form,

$$\mathcal{M} = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ s_2 & s_3 & s_4 & s_8 & s_7 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ s_2^2 & s_3^2 & s_4^2 & s_8^2 & s_7^2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ s_2^3 & s_3^3 & s_4^3 & s_8^3 & s_7^3 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & t_2 & t_3 & t_4 & t_8 & t_7 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & t_2 & t_3^2 & t_4^2 & t_8^2 & t_7^2 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & t_2 & t_3^3 & t_4^3 & t_8^3 & t_7^3 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & s_7 & s_6 & s_5 & s_1 & s_2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & s_7^2 & s_6^2 & s_5^2 & s_1^2 & s_2^2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & s_7^3 & s_6^3 & s_5^3 & s_1^3 & s_2^3 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ t_2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & t_7 & t_6 & t_5 \\ t_2^2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & t_7^2 & t_6^2 & t_5^2 \\ t_2^3 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & t_7^3 & t_6^3 & t_5^3 \end{pmatrix}$$

If $s_i = i$ and $t_j = j$, we can verify that the rank of the matrix is 15, thus the dimension of the spline space is 49. If we perturb one of the knots a little bit, such as $s_3 = 3.0 + \epsilon$, where ϵ is an arbitrary small value, then the dimension is 48. In other words, the dimension is instable.

5.2 Correction theorem

In this section, we will provide a new theorem to correct the theorem in [1] using the method in the last section.

Theorem 5.1. *Given a regular T-mesh $\mathcal{T}$ without holes, if each horizontal l-edges has at least $N^h - 1$ mono-vertices and each vertical l-edge has at least $N^v - 1$ mono-vertices except two end vertices, then the dimension of the spline space defined on $\mathcal{T}$ is*

$$\dim \mathcal{S}(d_1, d_2, \alpha, \beta, \mathcal{T}) = (d_1 + 1)(d_2 + 1) + (C^h - T^h)(d_1 + 1)(d_2 - \beta) + (C^v - T^v)(d_2 + 1)(d_1 - \alpha) + V(d_1 - \alpha)(d_2 - \beta).$$

Proof. We will prove that under the condition, the T-mesh is diagonalizable using reduction to absurdity. Actually, if the T-mesh is not diagonalizable. Then for any set of l-edges, it is bounded. Suppose e_i is the most bottom horizontal l-edges in the set (if there are more than one l-edge in the set, we




Figure 9: The new correction theorem for [1].

will pick the leftmost one), see Figure 9 as an illustration. According to the assumption, e_i has at least $N^h - 1$ mono-vertices (black rectangle vertices in the figure), if one of the two end vertices is not on the other l-edges of in the set, then e_i at least has N^h vertices which are not on the other l-edges in the set. According to Lemma 4.6, the T-mesh is diagonalizable which violates the assumption. Thus, there exists a vertical l-edge, e_j , which contains one of the end vertices of e_i . Without loss of generalization, we assume the right end vertex is on the l-edge e_j in the set. Now, we consider l-edge e_j , the bottom end vertex of the l-edge cannot lie on the other l-edges in the set because it is on the bottom of e_i which is the bottommost l-edges in the set. So e_j at least has N^v vertices which are not on the other l-edges in the set. According to Lemma 3.1, the T-mesh is diagonalizable which violates the assumption. Using Theorem 4.4, we prove the theorem. $\square$

6 Conclusion and Future work

In the present paper, we introduce a class of T-meshes, diagonalizable T-meshes, over which the dimension of spline spaces is stable. We also provide a necessary and sufficient condition to characterize this class of T-meshes. The dimension result in the present paper can cover all the existing dimension results as special cases.

The paper leaves several open problems for further research. As we have provided the dimension of the spline space, so there are many problems which need to be solved, such as construction of a set of basis functions with good properties, geometric operations and properties of the splines over T-meshes, etc. We will explore these problems in detail in future papers. It is also an important and interesting question to find out other general with fix dimension T-meshes.

Acknowledgements

The authors are supported by the NSF of China (No.11031007, No.60903148), Chinese Universities Scientific Fund, SRF for ROCS SE and Chinese Academy of Science (Startup Scientific Research Foundation).

References

- [1] C. J. Li, R. H. Wang, F. Zhang, Improvement on the Dimensions of Spline Spaces on T-Mesh, *Journal of Information & Computational Science* 3 (2) (2006) 235–244.
- [2] G. Farin, *NURBS Curves and Surfaces: from Projective Geometry to Practical Use*, Fourth Edition, A. K. Peters, Ltd., Natick, MA, 2002.
- [3] J. A. Cottrell, T. J. R. Hughes, Y. Bazilevs, *Isogeometric analysis: Toward Integration of CAD and FEA*, Wiley, Chichester, 2009.
- [4] J. Deng, F. Chen, Y. Feng, Dimensions of spline spaces over t-meshes, *Journal of Computational and Applied Mathematics* 194 (2006) 267–283.
- [5] J. Deng, F. Chen, X. Li, C. Hu, W. Tong, Z. Yang, Y. Feng, Polynomial splines over hierarchical t-meshes, *Graphical Models* 74 (2008) 76–86.
- [6] X. Li, J. Deng, F. Chen, Surface modeling with polynomial splines over hierarchical t-meshes, *The Visual Computer* 23 (2007) 1027–1033.
- [7] X. Li, J. Deng, F. Chen, Polynomial splines over general t-meshes, *The Visual Computer* 26 (2010) 277–286.
- [8] N. Nguyen-Thanh, H. Nguyen-Xuan, S. P. A. Bordas, T. Rabczuk, Isogeometric analysis using polynomial splines over hierarchical t-meshes for two-dimensional elastic solids, *Computer Methods in Applied Mechanics and Engineering* 200 (2011) 1892C1908.
- [9] P. Wang, J. Xu, J. Deng, F. Chen, Adaptive isogeometric analysis using rational pht-splines, *Computer-Aided Design* 43 (2011) 1438–1448.
- [10] L. Tian, F. Chen, Q. Du, Adaptive finite element methods for elliptic equations over hierarchical t-meshes, *J. Comput. Appl. Math.* 236 (2011) 878–891.
- [11] L. L. Schumaker, L. Wang, Splines on triangulations with hanging vertices, *Constructive Approximation* doi:10.1007/s00365-012-9167-x.
- [12] T. W. Sederberg, J. Zheng, A. Bakenov, A. Nasri, T-splines and T-NURCCs, *ACM Transactions on Graphics* 22 (3) (2003) 477–484.

- [13] T. W. Sederberg, D. L. Cardon, G. T. Finnigan, N. S. North, J. Zheng, T. Lyche, T-spline simplification and local refinement, *ACM Transactions on Graphics* 23 (3) (2004) 276–283.
- [14] T. W. Sederberg, G. T. Finnigan, X. Li, H. Lin, H. Ipson, Watertight trimmed NURBS, *ACM Transactions on Graphics* 27 (3) (2008) Article no. 79.
- [15] Y. Bazilevs, V. M. Calo, J. A. Cottrell, J. A. Evans, T. J. R. Hughes, S. Lipton, M. A. Scott, T. W. Sederberg, Isogeometric analysis using T-splines, *Computer Methods in Applied Mechanics and Engineering* 199 (5-8) (2010) 229 – 263.
- [16] Y. Bazilevs, L. Beirao de Veiga, J. Cottrell, T. Hughes, G. Sangalli, Isogeometric analysis: approximation, stability and error estimates for h -refined meshes, *Mathematical Models and Methods in Applied Sciences* 16 (2006) 1031–1090.
- [17] X. Li, J. Zheng, T. W. Sederberg, T. J. R. Hughes, M. A. Scott, On the linear independence of T-splines blending functions, *Computer Aided Geometric Design*, 29 (2012) 63–76.
- [18] M. A. Scott, X. Li, T. W. Sederberg, T. J. R. Hughes, Local refinement of analysis-suitable T-splines, *Computer Methods in Applied Mechanics and Engineering* 213-216 (2012) 206–222.
- [19] X. Li, M. A. Scott, Analysis-suitable t-splines: Characterization, refinability and approximation, submitted *Mathematical Models and Methods in Applied Sciences* for publish.
- [20] L. L. Schumaker, L. Wang, Spline spaces on tr-meshes with hanging vertices, *Numerische Mathematik* 118 (2011) 531–548.
- [21] B. Mourrain, On the dimension of spline spaces on planar t-subdivisions, *Arxiv preprint arXiv:1011.1752*.
- [22] R.-H. Wang, *Multivariate Spline Functions and Their Applications*, Science Press/ Kluwer Academic Publishers, 2001.
- [23] L. Schumaker, On the dimension of spaces of piecewise polynomials in two variables, in: *Multivariate Approximation Theory*, In: Schempp, W., Zeller, K. (Eds.), Birkhauser Verlag, Basel, 1979, pp. 396–412.
- [24] L. L. Schumaker, L. Wang, Approximation power of polynomial splines on t-meshes, *Computer Aided Geometric Design* doi:<http://dx.doi.org/10.1016/j.cagd.2012.04.003>.
- [25] A. Buffa, D. Cho, M. Kumar, Characterization of t-splines with reduced continuity order on t-meshes, *Comput. Methods Appl. Mech. Engrg.* 201-204 (2012) 112–126.

- [26] X. Li, F. Chen, On the instability in the dimension of spline space over particular t-meshes, *Computer Aided Geometric Design* 28 (2011) 420–426.
- [27] D. Berdinskya, M. jae Oha, T. wan Kima, B. Mourrain, On the problem of instability in the dimension of a spline space over a t-mesh, *Computers Graphics* 36(2) (2012) 507–513.
- [28] M. Wu, J. Deng, F. Chen, The dimension of spline spaces with highest order smoothness over hierarchical t-meshes, *Arxiv preprint arXiv:1112.1144*.